

X.—*Description of a new Species of Diurnal Lepidoptera belonging to the Genus Hebomoia.*—By J. WOOD-MASON.

The beautiful insect described below has been recently received by the Indian Museum from the Andaman Islands, where it was obtained by Mr. A. de Roepstorff, after whom I have named it.

HEBOMOIA ROEPSTORFII, n. sp.

♂. Differs from *H. glaucippe*, the only species of the genus with which I have been able to compare it, *on the upper side*, in having the apical orange patch of the fore-wing larger, extended into the cell, and less broadly bordered with black, both internally and externally; the submarginal black spots smaller and completely isolated from the black of the outer margin; the fore-wing at the posterior angle tinged, and the hind-wing externally broadly bordered, with bright sulphur-yellow, which colour is shaded off into the cream-colour of the rest of both wings; and the outer margin of the hind-wing narrowly edged with black, which gradually broadens from the anal to the anterior angle and extends inwards in points at the veins:—and, *on the under side*, in having the brown mottling of the fore-wing arranged in the form of a tolerably conspicuous band coincident with the macular band of the upper side; and the ground-colour of the hind-wing, as also that of the mottled portion of the fore-wing, of a rich golden-luteous colour.

Expanse 3·5 inches.

HAB. S. Andaman.

The place of this species would seem to be between *H. vossii*, (Maitland) and *H. sulphurea*, Wallace.

Presented  
8 MAY 1885